Information on Specific Tests and Vaccines

1. Tetanus, diphtheria and pertussis
   - Provide dates of your childhood vaccines
   - If you do not have documentation for your childhood immunizations (booklets with dates), you must be revaccinated following the adapted schedule as outlined on the immunization form
   - No booster dose is required before the age of 50 if the primary vaccination is completed
2. Polio
   - Systematic vaccination of adults, including healthcare workers interns, is not necessary in Quebec. The risk of exposure to the polio virus in North America is minimal.
   - Polio booster is not required unless the student is at increased risk of exposure to "wild" polio (e.g., those travelling to, or planning to work in areas that have wild polio or vaccine-derived polio outbreaks)
3. Measles

One of the following items is required:
   - Students born before 1970 are considered immune
   - Documented evidence of vaccination with two doses of measles containing vaccine, given at least a month a part starting on or after the first birthday
   - Documented history of the disease before January 1996
   - Documentation of positive measles serology** (attach results)
4. Mumps

One of the following items is required:
   - Students born before 1970 are considered immune
   - Documented evidence of vaccination with one dose of mumps containing vaccine
   - Documentation of positive mumps serology** (attach results)
5. Rubella

One of the following items is required:
   - Documented evidence of vaccination with one dose of Rubella containing vaccine
   - Documentation of positive Rubella serology** (attach results)

If serology results are negative to one or more of these 3 infections, a booster with a trivalent vaccine (MMR) is required.

Pre-test serological testing for measles, mumps and rubella is recommended, nor is the cost covered in Quebec. Students without written proof of vaccination should be vaccinated. Post testing following vaccination is also not recommended

6. Varicella

One of the following items is required:
   - A history of previous illness
   - Documented evidence of 2 doses of varicella vaccine (attach document)
   - Documentation of a positive varicella serology is required (attach document)

If you do not meet the above criteria, you must receive the vaccines (2 doses at least a month apart). No blood test is required after the vaccination. The CIQ recommends a total of 2 doses for interns and healthcare workers newly hired in the Quebec healthcare system.

7. Hepatitis B
   - Proof of vaccination: Complete series of 2 or 3 vaccines. If no documented proof of vaccination, proceed with the vaccination series. Serology before vaccination is not recommended nor it is covered in Quebec.
   - Serology is required if most recent dose was given in the last 1-6 months. It is not required if last dose of Hepatitis B vaccine was given more than 6 months ago.
   - In a case of accidental exposure to blood and/or body fluid a post exposure, evaluation is recommended.
   - Serology for Hepatitis B surface Antibody must be drawn 1 to 6 months after the last dose and show protective levels (anti-HBs H í ì Y Ł ¦ ĸ Y t ú ¢
   - If anti-HBs] • G í ì u / h/ml or absent, revaccinate with a single dose recheck for Hepatitis B surface Antibody and antigen 1 to 2 Months post vaccination. If anti- HBs H í ì u / h l u o v } (µ Œ š Z Œ
     If anti-HBs] • G í ì u / h Œ u o • v š } v š] v µ Á] š Z h š d Z •
     dose with the appropriate schedule AND recheck for Hepatitis B surface Antibody 1 to 2 months post vaccination.
   - Students found to have a chronic hepatitis B infection or to be a carrier (hepatitis B surface antigen positive - HBsAg) must consult their faculty's program coordinator.
   - Students found to be a non-converter after the second series of vaccines must consult their faculty's program coordinator.

8. Tuberculosis:

A two-step Mantoux is required ONLY for students answering YES to one of the following:

1. Born In Canada before 1976
2. Received the BCG vaccine
3. Known exposure/ contact with active tuberculosis
4. I as spent a cumulative period of 3 months or more in a country or countries with a high incidence of tuberculosis ó ĸ 3 л cases ĸ 1 л л Σ л л л inhabitants ĸ è Ŝ I · ₪ ú ¢ [λ ă ú 2 ₸ Ŏ 2 dz y ú ₪ Ŝ ă Ŏ I · y ¯ ō Ŝ ₸ 2 dz y Ř I · ú Ÿ K ú ú L J ă Ÿ ĸ ĸ ó ó ó ó ł y ă L J I j ¢ I j Ŏ ¢ Ŏ I · k ă I · y ú Ŝ π Ø 2 è I · Ǝ Ŝ ĸ Ǝ dz ŕ Ř Ŝ ĸ ₪ ă I j dz Ŝ ă k ú dz ō Ŝ ₪ Ŏ dz t 2 ă Ŝ k ă ĺ ú dz I · ú ĺ 2 y ¯

- All TST must be read 48-72 hours after administration by a trained healthcare worker. Self-reading of TST is not accepted
- The second TST is to be performed 7 days to 1 year after the first step provided there was no exposure to Mycobacterium tuberculosis during the past year.
- The q T Z Ø ê Â ' # ã s E ¼ ‰ ' ã ã ê Ã ì p d 1 0 6 9 T 8 < 0 3 6 >] T J / C 0 _ 0 9 T f

Student Wellness Hub
Student Services
3600 McTavish Street, Suite 3400 (West Wing)
Montreal, Quebec, Canada H3A 0G3

Pole bienêtre étudiant
Services aux étudiants
3600 rue McTavish, Bureau 3400 (Aile Ouest)
Montréal, Québec, Canada H3A 0G3

T : +1 514 398-6017
F : +1 514 398-2559